Application No.: 10/553,462 Docket No.: HO-P03236US0

AMENDMENTS TO THE CLAIMS

1. (Previously Presented) A method of identifying whether or not a pregnant woman is at risk of developing pre-eclampsia or whether or not her fetus is at risk of developing intrauterine growth restriction (IUGR), which method comprises:

- (a) measuring asymmetric dimethylarginine (ADMA) in a pregnant woman at a stage of pregnancy from 4 to 25 weeks gestation; and
- (b) determining whether or not the ADMA is greater than $2.0~\mu$ mol/L in the woman, thereby determining whether or not the woman is at risk of developing pre-eclampsia or her fetus is at risk of developing IUGR.
- 2. (Currently Amended) A <u>The</u> method according to <u>of</u> claim 1, wherein ADMA is measured in a fluid sample taken from the woman.
 - 3. (Cancel)
- 4. (Currently Amended) A <u>The</u> method according to <u>of</u> claim 1, wherein the pregnant woman is at a stage of pregnancy from 10 to 25 weeks gestation.
- 5. (Currently Amended) A <u>The</u> method according to <u>of</u> claim 4, wherein the woman is at a stage of pregnancy from 15 to 25 weeks gestation.
- 6. (Currently Amended) A The method according to of claim 1, wherein determining whether or not the woman is at risk of developing pre-eclampsia or determining whether or not her fetus is at risk of developing IUGR comprises determining whether or not the woman's ADMA level is at least 3 times the normal pregnancy level.
- 7. (Currently Amended) A <u>The</u> method according to <u>of</u> claim 1, wherein determining whether or not the woman is at risk of developing pre-eclampsia or determining whether or not her fetus is at risk of developing IUGR comprises determining whether or not

Application No.: 10/553,462 Docket No.: HO-P03236US0

the woman has an increase in the ADMA/symmetric dimethylarginine (ADMA/SDMA) ratio that is greater than the normal pregnancy ratio.

- 8. (Currently Amended) A <u>The</u> method according to <u>of</u> claim 7, comprising determining whether or not the ADMA/SDMA ratio is at least 5 times more than the normal pregnancy ratio.
- 9. (Currently Amended) A <u>The</u> method according to <u>of</u> claim 1, wherein the pregnant woman is suspected of being at risk of developing pre-eclampsia or her fetus is suspected of being at risk of developing IUGR.
- 10. (Currently Amended) A <u>The</u> method according to <u>of</u> claim 9, wherein the woman is a smoker.
- 11. (Currently Amended) A <u>The</u> method according to <u>of</u> claim 1, further comprising carrying out Doppler waveform analysis of the uterine arteries and/or flow-mediated dilatation of the brachial artery in the woman.

12.-17. (Canceled)

18. (Cancel)

19.-20. (Canceled)

21. (Cancel)

22. (Cancel)

23.-24. (Canceled)

25. (Cancel)

26. (Canceled)

27. (Cancel)

28. (Cancel) 60092161